



Corrigendum to 'Recovery of Oculomotor Nerve Palsy After Surgical and Endovascular Repair of Unruptured Internal Carotid-Posterior Communicating Artery Aneurysms': [World Neurosurg 188 (2024) e334– e340]

Shinya Tabata¹, Yushiro Take¹, Tatsuki Kimura¹, Koki Onodera¹, Haruka Kume¹, Masayoshi Fukuoka¹, Kaima Suzuki¹, Hidetoshi Ooigawa¹, Shinya Kohyama², Hiroki Kurita¹

■ The authors regret that in the original article, there were some mistakes in **Table 1**, which was published. Specifically, the notation “mean ± SD” following “age (years)” is incorrectly included and should be deleted, and the corresponding footnote “SD, standard deviation” is also unnecessary. “Diabetes” should be changed to “Diabetes mellitus”. Along with this, on page E335, in the “METHODS AND MATERIALS” section, “hypertension, diabetes, dyslipidemia” should be changed to “hypertension, diabetes mellitus, dyslipidemia”. The quartile range for

“time from onset to treatment (days)” in “All cases” should be changed to “5–47” instead of “570–47”. The corrected **Table 1** appears below. This oversight does not impact the study’s abstract, results, or conclusions.

The authors would like to apologise for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.wneu.2024.05.110>

Table 1. Baseline Characteristics of the Enrolled Patients With Oculomotor Nerve Palsy

	All Cases n=25	Clipping n=14	Coiling n=11	P Value
Age (years)	74 (61–78)	74 (62–77)	72 (62–82)	0.982
Female	23 (92.0)	13 (92.9)	10 (90.9)	1.000
Hypertension	16 (64.0)	8 (57.1)	8 (72.7)	0.677
Diabetes mellitus	6 (24.0)	3 (21.4)	3 (27.3)	1.000
Hyperlipidemia	12 (48.0)	4 (28.6)	8 (72.7)	0.047
Smoking	3 (12.0)	1 (7.1)	2 (18.2)	0.565
Ischemic heart disease	4 (16.0)	1 (7.1)	3 (27.3)	0.288
Complete ONP	9 (36.0)	6 (42.9)	3 (27.3)	0.677
Partial ONP	16 (64.0)	8 (57.1)	8 (72.7)	0.677
Symptom of Partial ONP				
Ptosis	9 (56.3)	5 (62.5)	4 (50.0)	1
Diplopia	14 (87.5)	7 (87.5)	7 (87.5)	1
Ophthalmoplegia	7 (43.8)	3 (37.5)	4 (50.0)	1
Impaired pupillary light reflex	3 (18.8)	2 (25.0)	1 (12.5)	1
Mydriasis	3 (18.8)	1 (12.5)	2 (25.0)	1
Aneurysm max diameter (mm)	6.4 (4.5–8.7)	6.2 (4.3–8.8)	6.3 (4.7–8.0)	0.979
Time from onset to treatment (days)	30 (5–47)	22 (8–70)	31 (5–39)	0.701
Data are presented as medians (interquartile range) or number (%).				
ONP, oculomotor nerve palsy.				

From the ¹Department of Cerebrovascular Surgery, International Medical Center, Saitama Medical University; and ²Department of Endovascular Neurosurgery, International Medical Center, Saitama Medical University, Hidaka, Japan

To whom correspondence should be addressed: Yushiro Take
[E-mail: y.take0804@gmail.com]

Citation: World Neurosurg. (2025).

<https://doi.org/10.1016/j.wneu.2024.12.001>

Journal homepage: www.journals.elsevier.com/world-neurosurgery

Available online: www.sciencedirect.com